

EPIDEMIOLOGY AND HERBAL TREATMENT OF OSTEOARTHRITIS

Syed Muhammad Ali Shah^{*1}, Muhammad Akram¹, Abid Rashid², Muhammad Daniyal³, Benish Ali⁴, Armghan Anjum⁴, Naheed Akhtar⁵, Imtiaz Mahmood Tahir⁵, Naveed Munir⁵

ABSTRACT Increase in prevalence of osteoarthritis has encouraged the researchers to find out novel, more efficacious drugs from plant sources. This study is a review of medicinal plants with anti-osteoarthritic effects in clinical trial and *in vitro/ in vivo* studies. Medicinal plants and herbal formulation/ products with anti-osteoarthritic effect are described. The plants and herbal products exerted anti-osteoarthritic effect by different mechanisms. The investigated plants contain the compounds that are able to contribute effectively to treat osteoarthritis. Therefore, the plants extract and herbal products discussed in this review article could open a way to conduct further clinical trials on large scales and greatly help researchers and pharmacists to develop new anti-osteoarthritic drugs.

1. Department of Eastern Medicine and Surgery, Directorate of Medical Sciences, Government College University Faisalabad, Pakistan
2. Directorate of Medical Sciences, Government College University Faisalabad, Pakistan
3. TCM and Ethnomedicine Innovation& Development Laboratory, School of Pharmacy, Hunan University of Chinese Medicine, Changsha, 410208, China.
4. Department of Orthotics and Prosthetics, Government College University Faisalabad Pakistan
5. College of Allied Health Professional, Directorate of Medical Sciences, Government College University Faisalabad, Pakistan

Corresponding author email: smalishah@hotmail.com

Key words: Arthritis, herbal treatment, efficacy

INTRODUCTION

Osteoarthritis is a common, inflammatory, skeletal, progressive, chronic and degenerative disorder (1). Protective cartilages on the ends of bones become worn, osteophytes form at the periphery of the joint surface and loose bodies may result (Fig. 1). Most common affected joints are knees and back (2-3). Manifestations of osteoarthritis include disability, swelling and pain in moving the joints (4). Disease progression can be slowed by proper treatment (Fig. 2). Despite ongoing physiotherapy and the widespread use of nonsteroidal anti-inflammatory drugs (NSAIDs), this illness is of apprehension because it affects the mobility of persons and life quality. Now there is bigger attention in non-synthetic, natural remedies obtained from plants to well acceptability and decreased adverse drug reactions (ADRs) (5-7). Plants have anti-arthritis activities (Table 1). Various medicinal plants have been proved as anti-arthritis agent in clinical trials (Table 2). In this review, we have reviewed literature regarding approved efficacy of plants in osteoarthritis.

Millions of people are affected with osteoarthritis. Its more prevalent in male than female before age of

45 and is more common in female than male after menopause. Its prevalence is increasing in United States and current 26 million people are affected with osteoarthritis in United State. Prevalence of osteoarthritis of distal interphalangeal joints was 75% in women between 60-70 years of age in United State. Prevalence of hand osteoarthritis in people above 40-50 years of age was 2.2% in Teheran that increased to 22.5% in people above 70 years of age. Study conducted by Dutch institute of public health indicated that prevalence of knee osteoarthritis is 15.6% in male and 30.5% in female. Prevalence is more in obese than normal weight persons. Prevalence of arthritis in Canadian adults was 13.4% and 17.6% in 1994 and 2002 respectively. Prevalence of arthritis was 16.9%, 14.2% and 14.7% in Ontario, Alberta and British Columbia. Prevalence of arthritis and rheumatism was 13.0%, 15.0%, 24%, 23.8% and 56.0% in the Medicinal plants having potential to treat arthritis. Medicinal plants have been traditionally used to control pain and improve dysfunction in osteoarthritis (Fig.3). In developing countries, most of people rely on herbal medicine (8-10). Herbal medicines are

prescribed worldwide for the management of osteoarthritis since ancient times (11). Herbs and plants always have been prescribed in the management of different diseases including gouty arthritis and other associated musculoskeletal disorders.

Zingiber officinale (Zingiberaceae).

This spice is a proven in nausea remedy (13). It can come in handy for the nausea associated with altitude sickness or provoked by driving on winding mountain roads (14). It is taken in various forms as a tea, tincture, capsules, raw or crystallized. It contains hexacosanoic acid 2,3-dihydroxypropyl ester, glycol monopalmitate, isovanillin, beta sitosterol palmitate, 1-(omega-ferulyloxygeranyl) glycerols, 6-shogaol, steroids, adenine, diterpenes, p-hydroxybenzaldehyde, gingerol analogue, maleimide-5-oxime and diarylheptanoids (15-17). It is anti-inflammatory, xanthine oxidase inhibitor, digestive, cyclooxygenase 2 inhibitor and antiarthritic (18). It is used in indigestion, inflammation, gouty arthritis and rheumatoid arthritis (19). Shimoda et al (2010) reported the anti-inflammatory effects of *Zingiber officinale* extract. This anti-inflammatory activity validates its efficacy in osteoarthritis (20).

A study was conducted to investigate the efficacy of *Zingiber officinale* extract as a substitute to NSAIDs and as a supplement medicine in the symptomatic management of osteoarthritis. The improvement of symptoms, defined as reduction in the mean change, was superior in the *Zingiber officinale* extract and ibuprofen groups than the placebo group. *Zingiber officinale* extract and ibuprofen were superior to placebo in the symptomatic treatment of osteoarthritis, while there was no significant difference between the *Zingiber officinale* extract and ibuprofen groups in a test for multiple comparisons (21).

Punica granatum (Lythraceae).

Chemical constituents include punicalagin, tannins and anthocyanins (22). It is used in pain, inflammation, arthritis, obesity, Alzheimer's disease, male infertility, infant brain ischemia, bacterial infections, erectile dysfunction, dental caries, diabetes mellitus, cardiovascular disorders

and cancer (23). Pharmacological activities are anticancer, diuretic, anti-angiogenesis, anti-mutagenic, astringent, antioxidant and anti-inflammatory (24). Pomegranate juice reduced proteoglycan loss and cartilage damage in mouse model with osteoarthritis (25).

Artemisia herba-alba Asso (Asteraceae)

Whole plant is prescribed for management of numerous disorders (26). It contains piperitone, carvone, cis-thujone, chrysanthemone and camphor (27). It contains trans-pinocarveol, camphor, borneol, alpha thujone, beta thujone, trans-sabinyl acetate, 1-8 cineole, chrysanthemone, cirsilineol and hispidulin (28). It is used in diabetes mellitus, osteoarthritis, nervous disorders, pyrexia, syphilis, scabies, neuralgia, diarrhea, bronchitis, cough, cold and fungal infection. It is anti-inflammatory, antioxidant, hypoglycemic, antileishmania and smooth muscle relaxant (29). Arshad et al (2012) reported the ethno medicinal use of this plant in arthritis (30).

Harpagophytum procumbens (Pedaliaceae)

This plant is used to treat various ailments (31). This plant contains iridoids that possess anti-inflammatory effect. Chantre et al (2000) reported the potential of *Harpagophytum procumbens* in comparison with diacerhein in management of osteoarthritis in randomized clinical study. This was a comparative study, in which efficacy of herbal product (Harpadol) was investigated in comparison with diacerhein. Six capsules of herbal medicine were given to patient daily. Each capsule contains 435 mg of powdered material. Diacerhein was given at dose of 100 mg/day. The duration of treatment was four months. Total no of patients with osteoarthritis was 122. Pain was significantly reduced in both management clusters. Efficacy of both drugs was comparable. After completion of trial duration, Harpadol treated patients were taking significantly less NSAIDs and analgesic drugs. Fewer side effects occurred in harpadol group than control group. 8.1% patients were suffered from diarrhea and 26.7% were in control group. This research indicated that test drug is equal in effect and better in safety to control group (32).

Dalbergia sissoo Roxb (Fabaceae)

Parts used are roots and wood. Active principles are isoflavones, flavonols and lignan glucoside (33). It is used in arthritis, boils, eruption of skin, leprosy, vomiting and inflammation (34). It is astringent, anti-inflammatory, anti-arthritic and gastroprotective (35). Dixit et al (2010) reported the ingredients from leaves of *Dalbergia sissoo* possessing estrogenic effect. Ethanol extract of this plant contains various constituents that have osteogenic effect in primary calvarial osteoblast cultures. Some compounds of this plant increased alkaline phosphatase activity. Osteogenic activity is evident from mineralization (36).

Commiphora mukul (Burseraceae)

It is used in hemorrhoids (37). It is laxative, antiseptic, anti-catarhal, expectorant, carminative, astringent, antimicrobial, anti-inflammatory, demulcent, hepatoprotective and anti-diabetic (38). Singh et al (2003) reported the effectiveness of *Commiphora mukul* for osteoarthritis of the knee. *Commiphora mukul* has been investigated for its efficacy to treat osteoarthritis. Preclinical and clinical data shows that *Commiphora mukul* is effective in osteoarthritis. In a clinical trial, thirty male and female patients were selected for study. *Commiphora mukul* was administered to patients at dose of 500 mg in form of capsule. There was significant improvement in mobility and functions of joints when taken for 1 month continuously. Treatment was continued to improve at the 2-month marker and follow-up. *Commiphora mukul* was effective in participants during the trial. During the trial, no significant adverse effects were reported. This study showed that *Commiphora mukul* is safe and effective alternative to treat osteoarthritis (39).

Curcuma longa (Zingiberaceae).

It is used in rheumatism. It contains curcumin, bisdemethoxycurcumin, demethoxycurcumin, zingiberone, tumerone, atlantone, resins, protein and sugars (40). It is used in acute coronary syndrome, psoriasis, vitiligo, arthritis, cardiovascular disorders, tumor, gastric ulcer, peptic ulcer, diabetes mellitus, ulcerative proctitis,

gastric inflammation, crohn's disease, diabetic nephropathy, ulcerative colitis, lupus nephritis, irritable bowel disease, acquired immune deficiency syndrome, tropical pancreatitis, atherosclerosis, idiopathic orbital inflammatory pseudotumor, diabetic microangiopathy, oral lichen planus, β -thalassemia, chronic bacterial prostatitis, biliary dyskinesia and cholecystitis (41). It is anti-inflammatory, antioxidant and anti-cancer (42). Cucumin containing formulation exhibited anti-inflammatory activity in patients with osteoarthritis (43).

Boswellia serrata (Burseraceae)

Chemical constituents are pentacyclic triterpene acids and beta-boswellic acid (44). It is used in asthma, inflammation, cancer and osteoarthritis (45). Pharmacological activities are anticancer, antioxidant, anti-asthmatic and anti-inflammatory (46). *Boswellia serrata* is prescribed in the treatment of osteoarthritis (47). Kimmatkar et al (2003) reported the efficacy of *Boswellia serrata* extract in management of knee osteoarthritis. In a clinical trial, 56 patients were randomized into two groups. 500 mg capsule of *Boswellia serrata* was administered to first group in three divided doses. Capsules were administered with lukewarm water. Total patients were 29 in first group. 23 patients were in 2nd group and capsule was administered with Luke water. In 2nd group, *Boswellia serrata* ointment was applied on joints. This treatment was given for 2 month duration. There was symptomatic improvement in both groups. Improvement in first group was with promising results (48).

Phyllanthus emblica (Euphorbiaceae)

Fruit of this plant is used in osteoarthritis (49). Chemical constituents include minerals, amino acids, emblicol, curcuminoids, phyllembelic acid, tannins and phenolic compounds (50). It is used in inflammation, jaundice, diarrhea and cancer (51). Pharmacological activities include antioxidant, gastroprotective, hepatoprotective, antiulcerogenic, antibacterial, hypolipidemic and anticancer (52). Sumantran et al (2008) reported the chondroprotective activity of *Phyllanthus emblica* in patients with osteoarthritis. Aqueous

extract of *Phyllanthus emblica* was investigated for its chondroprotective activity. In *In vitro* study, hyaluronidase and collagenase type

2 activities were inhibited by use of aqueous extract of *Phyllanthus emblica*. This data shows that *Phyllanthus emblica* extract can be used as chondroprotective agent in the treatment of osteoarthritis (53).

CONCLUSION

The current review reveals the anti-osteoarthritis efficacies of commonly used medicinal plants in Unani system of medicine. Since natural medicines are worldwide prescribed and the bio-active constituents isolated from these herbs may function as bio-marker in different natural formulations. Present review also supports the use of these herbal preparations that are used to prevent or control the inflammatory conditions.

REFERENCES

1. Zhou Y, Wang T, Hamilton JL, Chen D. Wnt/ β -catenin signaling in osteoarthritis and in other forms of arthritis. *Current rheumatology reports*. 2017;19(9):53.
2. Yudoh K, Terauchi K, Yui N, Kobayashi H, Fujiya H, Niki H, et al. Impact of NAD-dependent deacetylase Sirtuin-1 in the osteophyte formation and the degradation of articular cartilage in osteoarthritis (OA). *Osteoarthritis and Cartilage*. 2017;25:S152-S3.
3. Clegg DO, Reda DJ, Harris CL, Klein MA, O'dell JR, Hooper MM, et al. Glucosamine, chondroitin sulfate, and the two in combination for painful knee osteoarthritis. *New England Journal of Medicine*. 2006;354(8):795-808.
4. Felson DT, Zhang Y. An update on the epidemiology of knee and hip osteoarthritis with a view to prevention. *Arthritis & Rheumatology*. 1998;41(8):1343-55.
5. Brahmachari B, Chatterjee S, Ghosh A. Efficacy and safety of diacerein in early knee osteoarthritis: a randomized placebo-controlled trial. *Clinical rheumatology*. 2009;28(10):1193-8.
6. Cisár P, Jány R, Waczulíková I, Sumegová K, Muchová J, Vojtaššák J, et al. Effect of pine bark extract (Pycnogenol®) on symptoms of knee osteoarthritis. *Phytotherapy Research*. 2008;22(8):1087-92.
7. Colker CM, Swain M, Lynch L, Gingerich DA. Effects of a milk-based bioactive micronutrient beverage on pain symptoms and activity of adults with osteoarthritis: a
8. double-blind, placebo-controlled clinical evaluation. *Nutrition*. 2002;18(5):388-92.
9. Karimifar M, Soltani R, Hajhashemi V, Sarrafchi S. Evaluation of the effect of *Elaeagnus angustifolia* alone and combined with *Boswellia thurifera* compared with ibuprofen in patients with knee osteoarthritis: a randomized double-blind controlled clinical trial. *Clinical rheumatology*. 2017;36(8):1849-53.
10. Chopra A, Lavin P, Patwardhan B, Chitre D. A 32-week randomized, placebo-controlled clinical evaluation of RA-11, an Ayurvedic drug, on osteoarthritis of the knees. *JCR: Journal of Clinical Rheumatology*. 2004;10(5):236-45.
11. Hoareau L, DaSilva EJ. Medicinal plants: a re-emerging health aid. *Electronic Journal of biotechnology*. 1999;2(2):3-4.
12. Sen S, Chakraborty R. Revival, modernization and integration of Indian traditional herbal medicine in clinical practice: Importance, challenges and future. *Journal of traditional and complementary medicine*. 2017;7(2):234-44.
13. Nirmal P, Koppikar S, Bhondave P, Narkhede A, Nagarkar B, Kulkarni V, et al. Influence of six medicinal herbs on collagenase-induced osteoarthritis in rats. *The American journal of Chinese medicine*. 2013;41(06):1407-25.
14. Ryan JL, Heckler CE, Roscoe JA, Dakhil SR, Kirshner J, Flynn PJ, et al. Ginger (*Zingiber officinale*) reduces acute chemotherapy-induced nausea: a URCC CCOP study of 576 patients. *Supportive care in cancer*. 2012;20(7):1479-89.
15. Akram M, Shah MI, Usmanghan K,

- Mohiuddin E, Sami A, Asif M, et al. Zingiber officinale roscoe (A medicinal plant). Pakistan Journal of Nutrition. 2011;10(4):399- 400.
16. Ali BH, Blunden G, Tanira MO, Nemmar A. Some phytochemical, pharmacological and toxicological properties of ginger (Zingiber officinale Roscoe): a review of recent research. Food and chemical Toxicology. 2008;46(2):409-20.
17. Jolad SD, Lantz RC, Solyom AM, Chen GJ, Bates RB, Timmermann BN. Fresh organically grown ginger (Zingiber officinale): composition and effects on LPS-induced PGE2 production. Phytochemistry. 2004;65(13):1937-54.
18. Jolad SD, Lantz RC, Chen GJ, Bates RB, Timmermann BN. Commercially processed dry ginger (Zingiber officinale): composition and effects on LPS-stimulated PGE2 production. Phytochemistry. 2005;66(13):1614- 35.
19. Kashefi F, Khajehei M, Alavinia M, Golmakani E, Asili J. Effect of Ginger (Zingiber officinale) on Heavy Menstrual Bleeding: A Placebo Controlled, Randomized Clinical Trial. Phytotherapy Research. 2015;29(1):114-9.
20. Grzanna R, Lindmark L, Frondoza CG. Ginger—an herbal medicinal product with broad anti-inflammatory actions. Journal of medicinal food. 2005;8(2):125-32.
21. Shimoda H, Shan S-J, Tanaka J, Seki A, Seo J-W, Kasajima N, et al. Anti-inflammatory properties of red ginger (Zingiber officinale var. Rubra) extract and suppression of nitric oxide production by its constituents. Journal of medicinal food. 2010;13(1):156-62.
22. Haghighi M, Khalvat A, Toliat T, Jallaei S. Comparing the effects of ginger (Zingiber officinale) extract and ibuprofen on patients with osteoarthritis. 2005.
23. Wang R-F, Xie W-D, Zhang Z, Xing D-M, Ding Y, Wang W, et al. Bioactive Compounds from the Seeds of Punica granatum (Pomegranate). Journal of Natural Products. 2004;67(12):2096-8.
24. Jurenka J. Therapeutic applications of pomegranate (Punica granatum L.): a review. Alternative medicine review. 2008;13(2):128.
25. Rahimi HR, Arastoo M, Ostad SN. A comprehensive review of Punica granatum (pomegranate) properties in toxicological, pharmacological, cellular and molecular biology researches. Iranian journal of pharmaceutical research: IJPR. 2012;11(2):385.
26. Hadipour Jahromy M, Mozaffari Kermani R. Chondroprotective effects of pomegranate juice on monoiodoacetate induced osteoarthritis of the knee joint of mice. Phytotherapy Research. 2010;24(2):182-5.
27. Abad MJ, Bedoya LM, Apaza L, Bermejo P. The Artemisia L. genus: a review of bioactive essential oils. Molecules. 2012;17(3):2542-66.
28. Amri I, De Martino L, Marandino A, Lamia H, Mohsen H, Scandolera E, et al. Chemical composition and biological activities of the essential oil from Artemisia herba-alba growing wild in Tunisia. Nat Prod Comm. 2013;8:407- 10.
29. Salah SM, Jäger AK. Two flavonoids from Artemisia herba-alba Asso with in vitro GABAA-benzodiazepine receptor activity. Journal of Ethnopharmacology. 2005;99(1):145-6.
30. Aziz M, Karim A, EL OUARIACHI EM, BOUYANZER A, AMRANI S, MEKHFI H, et al. Relaxant effect of essential oil of Artemisia herba-alba Asso. on rodent jejunum contractions. Scientia pharmaceutica. 2012;80(2):457-68.
31. Shedayi AA, Gulshan B. Ethnomedicinal uses of plant resources in Gilgit-Baltistan of Pakistan. Journal of Medicinal Plants Research. 2012;6(29):4540-9.
32. Kundu JK, Mossanda KS, Na H-K, Surh Y-J. Inhibitory effects of the extracts of Sutherlandia frutescens (L.) R. Br. and Harpagophytum procumbens DC. on phorbol ester-induced COX-2 expression in mouse skin: AP-1 and CREB as potential upstream targets. Cancer letters. 2005;218(1):21-31.
33. Chantre P, Cappelaere A, Leblan D, Guedon D, Vandermander J, Fournie B. Efficacy and tolerance of Harpagophytum

- procumbens versus diacerhein in treatment of osteoarthritis. *Phytomedicine*. 2000;7(3):177-83.
34. Asif M, Kumar A. Phytochemical investigation and evaluation of antinociceptive activity of ethanolic extract of *Dalbergia sissoo* (Roxb.) bark. *Journal of natural science, biology, and medicine*. 2011;2(1):76.
35. Hajare S, Chandra S, Sharma J, Tandan S, Lal J, Telang a. Anti-inflammatory activity of *Dalbergia sissoo* leaves. *Fitoterapia*. 2001;72(2):131-9.
36. Khan MI, Khan MR. Gastroprotective potential of *Dalbergia sissoo* roxb. stem bark against diclofenac- induced gastric damage in rats. *Osong public health and research perspectives*. 2013;4(5):271-7.
37. Dixit P, Chillara R, Khedgikar V, Gautam J, Kushwaha P, Kumar A, et al. Constituents of *Dalbergia sissoo* Roxb. leaves with osteogenic activity. *Bioorganic & medicinal chemistry letters*. 2012;22(2):890-7.
38. Yousefi M, Mahdavi MRV, Hosseini SM, Bahrami A, Davati A, Kamalinejad M, et al. Clinical Evaluation of Commiphora Mukul, a Botanical resin, in the Management of Hemorrhoids: A randomized controlled trial. *Pharmacognosy magazine*. 2013;9(36):350.
39. Ramesh B, Karuna R, Reddy SS, Sudhakara G, Saralakumari D. Ethanolic extract of Commiphora mukul gum resin attenuates streptozotocin-induced alterations in carbohydrate and lipid metabolism in rats. *EXCLI journal*. 2013; 12:556.
40. Singh BB, Mishra LC, Vinjamury SP, Aquilina N. The effectiveness of Commiphora mukul for osteoarthritis of the knee: an outcomes study. *Alternative therapies in health and medicine*. 2003;9(3):74.
41. Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: a review of preclinical and clinical research. *Alternative medicine review*. 2009;14(2).
42. Gupta SC, Patchva S, Aggarwal BB. Therapeutic roles of curcumin: lessons learned from clinical trials. *The AAPS journal*. 2013;15(1):195-218.
43. Kocaadam B, Şanlıer N. Curcumin, an active component of turmeric (*Curcuma longa*), and its effects on health. *Critical reviews in food science and nutrition*. 2017;57(13):2889-95.
44. Akram M, Shahab-Uddin AA, Usmanghani K, Hannan A, Mohiuddin E, Asif M. Curcuma longa and curcumin: a review article. *Romanian Journal of Plant Biology*. 2010;55(2):65-70.
45. Ahmed HH, Abd-Rabou AA, Hassan AZ, Kotob SE. Phytochemical analysis and anti-cancer investigation of *Boswellia serrata* bioactive constituents in vitro. *Asian Pacific Journal of Cancer Prevention*. 2015;16(16):7179-88.
46. Gupta A, Khajuria A, Singh J, Singh S, Suri K, Qazi G. Immunological adjuvant effect of *Boswellia serrata* (BOS 2000) on specific antibody and cellular response to ovalbumin in mice. *International immunopharmacology*. 2011;11(8):968-75.
47. Siddiqui M. *Boswellia serrata*, a potential antiinflammatory agent: an overview. *Indian journal of pharmaceutical sciences*. 2011;73(3):255.
48. Kimmatkar N, Thawani V, Hingorani L, Khiyani R. Efficacy and tolerability of *Boswellia serrata* extract in treatment of osteoarthritis of knee-a randomized double blind placebo controlled trial. *Phytomedicine*. 2003;10(1):3-7.
49. Gupta P, Samarakoon S, Chandola H, Ravishankar B. Clinical evaluation of *Boswellia serrata* (Shallaki) resin in the management of Sandhivata (osteoarthritis). *Ayurvedic*. 2011;32(4):478.
50. Ram MS, Neetu D, Yogesh B, Anju B, Dipti P, Pauline T, et al. Cyto-protective and immunomodulating properties of Amla (*Emblica officinalis*) on lymphocytes: an in-vitro study. *Journal of Ethnopharmacology*. 2002;81(1):5-10.
51. Gaire BP, Subedi L. Phytochemistry, pharmacology and medicinal properties of *Phyllanthus emblica* Linn. *Chinese journal of integrative medicine*. 2014:1-8.
52. Baliga MS, Dsouza JJ. Amla (*Emblica officinalis* Gaertn), a wonder berry in the treatment and prevention of cancer. *European Journal of*

- Cancer Prevention. 2011;20(3):225-39.
53. Krishnaveni M, Mirunalini S, Karthishwaran K, Dhamodharan G. Antidiabetic and antihyperlipidemic properties of *Phyllanthus emblica* Linn.(Euphorbiaceae) on streptozotocin induced diabetic rats. *Pakistan Journal of Nutrition*. 2010;9(1):43-51.
54. Sumantran VN, Kulkarni A, Chandwaskar R, Harsulkar A, Patwardhan B, Chopra A, et al. Chondroprotective potential of fruit extracts of *Phyllanthus emblica* in osteoarthritis. *Evidence-Based Complementary and Alternative Medicine*. 2008;5(3):329-35.
55. Patwardhan SK, Bodas KS, Gundewar S. Coping with arthritis using safer herbal options. *International Journal of Pharmacology*. 2010;2(1):1-11.
56. Bansod MS, Kagathara V, Pujari R, Patel V, Ardeshta H. Therapeutic effect of a poly-herbal preparation on adjuvant induced arthritis in wistar rats. *International journal of pharmacy and pharmaceutical sciences*. 2011;3(2):186- 92.
57. Yesilada E, Küpeli E. *Clematis vitalba* L. aerial part exhibits potent anti-inflammatory, antinociceptive and antipyretic effects. *Journal of Ethnopharmacology*. 2007;110(3):504-15.
- Andersen ML, Santos EH, Maria de Lourdes VS, da Silva AA, Tufik S. Evaluation of acute and chronic treatments with *Harpagophytum procumbens* on Freund's adjuvant-induced arthritis in rats. *Journal of Ethnopharmacology*. 2004;91(2-3):325-30.
58. Jung H-J, Nam JH, Choi J, Lee K-T, Park H-J. Antiinflammatory effects of chiisanoside and chiisanogenin obtained from the leaves of *Acanthopanax chiisanensis* in the carrageenan- and Freund's complete adjuvant- induced rats. *Journal of Ethnopharmacology*. 2005;97(2):359-67.
59. Amresh G, Singh P, Rao CV. Antinociceptive and antiarthritic activity of *Cissampelos pareira* roots. *Journal of Ethnopharmacology*. 2007;111(3):531-6.
57. Kim K-S, Lee S-D, Kim K-H, Kil S-Y, Chung K-H, Kim C-H. Suppressive effects of a water extract of *Ulmus davidiana* Planch (Ulmaceae) on collagen-induced arthritis in mice. *Journal of Ethnopharmacology*. 2005;97(1):65-71.
58. Ojewole JA. Evaluation of the anti-inflammatory properties of *Sclerocarya birrea* (A. Rich.) Hochst. (family: Anacardiaceae) stem-bark extracts in rats. *Journal of Ethnopharmacology*. 2003;85(2-3):217-20.
59. Fan A, Lao L, Zhang R, Zhou A, Wang L, Moudgil K, et al. Effects of an acetone extract of *Boswellia carterii* Birdw.(Burseraceae) gum resin on adjuvant-induced arthritis in lewis rats. *Journal of Ethnopharmacology*. 2005;101(1- 3):104-9.
60. Perez R, Perez S, Zavala M, Salazar M. Anti-inflammatory activity of the bark of *Hippocratea excelsa*. *Journal of Ethnopharmacology*. 1995;47(2):85-90.
61. Olajide OA, Makinde JM, Okpako DT, Awe SO. Studies on the anti-inflammatory and related pharmacological properties of the aqueous extract of *Bridelia ferrugineastem* bark. *Journal of Ethnopharmacology*. 2000;71(1-2):153-60.
62. Choi J, Huh K, Kim S-H, Lee K-T, Lee H-K, Park H-J. Kalopanaxsaponin A from *Kalopanax pictus*, a potent antioxidant in the rheumatoid rat treated with Freund's complete adjuvant reagent. *Journal of Ethnopharmacology*. 2002;79(1):113-8.
63. Omisore N, Adewunmi C, Iwalewa E, Ngadju B, Watchueng J, Abegaz B, et al. Antinociceptive and anti-inflammatory effects of *Dorstenia barteri* (Moraceae) leaf and twig extracts in mice. *Journal of Ethnopharmacology*. 2004;95(1):7-12.
64. Ojewole JA, Adewunmi CO. Anti-inflammatory and hypoglycaemic effects of *Tetrapleura tetraptera* (Taub) [fabaceae] fruit aqueous extract in rats. *Journal of Ethnopharmacology*. 2004;95(2-3):177-82.
65. Kuptniratsaikul V, Thanakhumtorn S, Chinswangwatanakul P, Wattanamongkonsil L, Thamlikitkul V. Efficacy and safety of *Curcuma domestica* extracts in patients with knee osteoarthritis. *The Journal of Alternative and Complementary Medicine*. 2009;15(8):891-7.

68. Panahi Y, Alishiri GH, Parvin S, Sahebkar A. Mitigation of systemic oxidative stress by curcuminoids in osteoarthritis: results of a randomized controlled trial. *Journal of dietary supplements*. 2016;13(2):209-20.
69. Panahi Y, Alishiri GH, Bayat N, Hosseini SM, Sahebkar a. Efficacy of *Elaeagnus Angustifolia* extract in the treatment of knee osteoarthritis: a randomized controlled trial. *EXCLI journal*. 2016; 15:203.
70. Soleimani H, Kamalinejad M, Kor NM, Najafizadeh SR, Askarfarashah M, HoseiniYekta N, et al. A preliminary study on the efficacy of *Nerium oleander* L. extract on clinical symptoms in patients with knee osteoarthritis. *European Journal of Experimental Biology*. 2014;4(2):295-8.
71. Lecomte A, Costa J. *Harpagophytum* dans l'arthrose: Etude en double insu contre placebo. *Le Magazine*. 1992; 15:27-30.
72. Randall C, Randall H, Dobbs F, Hutton C, Sanders H. Randomized controlled trial of nettle sting for treatment of base-of-thumb pain. *Journal of the Royal Society of Medicine*. 2000;93(6):305- 9.
73. Schmid B, Tschirdewahn B, Kötter I, Günaydin I, Lüdtke R, Selbmann H, et al. Analgesic effects of willow bark extract in osteoarthritis: results of a clinical double blind trial. *Focus on Alternative and Complementary Therapies*. 1998;3(4):186-191.
74. Kulkarni R, Patki P, Jog V, Gandage S, Patwardhan B. Treatment of osteoarthritis with a herbomineral formulation: a double-blind, placebo-controlled, cross-over study. *Journal of Ethnopharmacology*. 1991;33(1-2):91-5.
75. Maheu E, Mazières B, Valat JP, Loyau G, Loët XL, Bourgeois P, et al. Symptomatic efficacy of avocado/ soybean unsaponifiables in the treatment of osteoarthritis of the knee and hip: A prospective, randomized, double blind, placebo controlled, multicenter clinical trial with a six-month treatment period and a two month followup demonstrating a persistent effect. *Arthritis & Rheumatology*. 1998;41(1):81-91.
76. Altman RD, Aven A, Holmburg CE, Pfeifer LM, Sack M, Young GT, editors. Capsaicin cream 0.025% as monotherapy for osteoarthritis: a double-blind study. *Seminars in arthritis and rheumatism*; 1994: Elsevier.
77. Biswas N. Treatment of osteoarthritis, rheumatoid arthritis and nonspecific arthritis with a herbal drug: a double-blind, active drug controlled parallel study. *JK Pract*. 1998; 5:129-32.
78. Rytting K, Schlamowitz P, Warnøe O, Wilstrup F. Gitadyl versus ibuprofen in patients with osteoarthrosis. The result of a double-blind, randomized cross-over study. *Ugeskrift for laeger*. 1991;153(33):2298-9.

Table 1: Plants investigated for their anti-arthritic activity in animal model

S. No	Plant Name	Origin of plant	Use of plant	Use in which part of the World	Studies in animals	References
1	<i>Punica granatum</i>	Iran, India and Pakistan	Antioxidant and anti-inflammatory.	Asia, Iran, Africa, India and Europe	Mouse model	(25)
2	<i>Semecarpus anacardium</i>	India	Anti-inflammatory, antiarthritic	India	Freund's adjuvant induced arthritis	(54)
3	<i>Withania somnifera</i>	Europe and Africa	Anti-inflammatory, antiarthritic	Pakistan and India	Carrageenan induced paw edema	(54)
4	<i>Vitex negunda</i>	Bhutan, Bangladesh, Afghanistan	Antiinflammatory, antiarthritic	Pakistan and India	Freud adjuvant induced arthritis	(55)
5	<i>Ricinus communis</i>	Northeastern Africa and Middle East	Contraceptive and antiarthritic	Malta, Pakistan and India	Freund's adjuvant induced arthritis	(54)
6	<i>Clematis vitalba</i>	England	Antimicrobial, Antiarthritic	Italy, Turkey	Carrageenan induced paw edema	(56)
7	<i>Harpagophytum procumbens</i>	Africa	Analgesic, Antiarthritic	United Kingdom, Scotland	Freund's adjuvant induced arthritis	(57)
8	<i>Acanthopanax chiisanensis</i>	South Asia	Antiinflammatory, Antiarthritic	Korea, Japan	Freund's adjuvant induced arthritis	(58)
9	<i>Cissampelos pareira</i>	Nepal	Anticancer, Antiarthritic	India, Bangladesh	Freud adjuvant induced arthritis	(59)
10	<i>Ulmus davidiana</i>	Japan, Siberia, Mongolia, and Korea	Anti-inflammatory, Antiarthritic	Korea	Collagen induced arthritis	(60)

11	<i>Sclerocarya birrea</i>	Africa	Antibacterial, Antiarthritic	Africa, Zimbabwe	Carrageen an induced paw edema	(61)
12	<i>Boswellia carteri</i>	Africa, Yemen and Oman	Anticancer, Antiarthritic	Egypt, Japan	Adjuvant induced arthritis	(62)
13	<i>Hippocratea excels</i>	México and Central America	Gastroprotective, Antiarthritic	Mexico, Japan	Adjuvant induced arthritis	(63)
14	<i>Bridelia ferruginea</i>	South Asia and Africa	Antiinflammatory, Antiarthritic	Nigeria and Germany	Adjuvant induced arthritis	(64)
15	<i>Kalopanax pictus</i>	China, Korea and Japan	Antiinflammatory, Antiarthritic	Korea	Freund's adjuvant induced arthritis	(65)
16	<i>Dorstenia barteri</i>	Cameroun and Nigeria	Antiinflammatory, Antiarthritic	Nigeria, France	Carrageen an induced paw edema	(66)
17	<i>Tetrapleura tetraaptera</i>	Western Africa	Antiinflammatory, Antiarthritic	South Africa, Nigeria	Egg albumin induced paw edema	(67)

Table 2: Clinical trials conducted for investigation of plants and herbal products in patients with arthritis

S. No	Plant Name/Formulation	Use of plant	No of Patients	References
1	<i>Curcuma domestica</i>	Antioxidant, anti-Inflammatory	52	(68)
2	<i>Boswellia serrata</i>	Analgesic, antiarthritic and anti-inflammatory	15	(47)
3	<i>Zingiber officinale</i>	Antiinflammatory and Analgesic	40	(21)
4	<i>Elaeagnus angustifolia</i>	Wound healer, muscle relaxant, antiinflamamatory	97	(69-70)
5	<i>Nerium oleander</i>	Anti-inflammatory	18	(71)
6	<i>Harpagophytum procumbens</i>	Anti-inflammatory	89	(72)
7	Stinging nettle	Anti-inflammatory	27	(73)
8	Willow bark	Anti-inflammatory	78	(74)
9	Articulín-F	Anti-inflammatory	42	(75)
10	Avocado/soybean unsaponifiables	Anti-inflammatory	164	(76)
11	Capsaicin	Anti-inflammatory	133	(77)
12	Eazmov	Anti-inflammatory	60	(78)
13	Gitadyl	Anti-inflammatory	35	(79)
14	Reumalex	Anti-inflammatory	82	(79)

Fig 1: Molecular mechanism underlying pathophysiology of osteoarthritis




Fig 3: Medicinal plants having potential to treat osteoarthritis

